

A Note on the Asymptotic Limit of the Four Simplex

Suresh. K. Maran

February 7, 2008

Abstract

Recently the asymptotic limit of the Barrett-Crane models has been studied by Barrett and Steele. Here by a direct study, I show that we can extract the bivectors which satisfy the essential Barrett-Crane constraints from the asymptotic limit. Because of this the Schläfli identity is implied by the asymptotic limit, rather than to be imposed as a constraint.

The asymptotic limit [7], [6], [4] of the Barrett-Crane models [2], [3] has been recently studied systematically in [5]. Here by a direct study, I show that we can extract the bivectors which satisfy the essential Barrett-Crane constraints¹ from the asymptotic limit. Because of this the Schläfli identity is implied by the asymptotic limit, rather than to be imposed as a constraint as in Ref:[2]. Here I focus on the Riemannian Barrett-Crane model [2] only, but it can be generalized to the Lorentzian Barrett-Crane model [3].

Consider the amplitude of a four-simplex[2] with a real scale parameter λ ,

$$\begin{aligned} Z_\lambda &= (-1)^{\sum_{k<l} 2J_{kl}} \int_{n_k \in S^3} \prod_{i<j} \frac{\sin(\lambda J_{ij} \theta_{ij})}{\sin(\theta_{ij})} \prod_k dn_k, \\ &= \frac{(-1)^{\sum_{k<l} 2J_{kl}}}{(2i)^{10}} \int_{n_k \in S^3} \prod_{i<j} \sum_{\varepsilon_{ij}=\pm 1} \frac{\varepsilon_{ij} \exp(i\varepsilon_{ij} \lambda (\rho J_{ij} \theta_{ij}))}{\sin(\theta_{ij})} \prod_k dn_k, \end{aligned}$$

where the θ_{ij} is defined by $n_i \cdot n_j = \cos(\theta_{ij})$. Here the θ_{ij} is the angle between n_i and n_j . The asymptotic limit of $Z_\lambda(s)$ under $\lambda \rightarrow \infty$ is controlled by

$$\begin{aligned} S(\{n_i\}, \{J_{ij}\}, \{q_i\}) \\ = \sum_{i<j} \varepsilon_{ij} J_{ij} \theta_{ij} + \sum_i q_i (n_i \cdot n_i - 1), \end{aligned}$$

where the q_i are the Lagrange multipliers to impose $n_i \cdot n_i = 1, \forall i$. My goal now is to find stationary points for this action. The stationary value under the

¹Here by the essential Barrett-Crane constraints [2] I mean all the constraints except those that relate to degeneracy conditions of a four simplex.

variation of n_j 's are determined by

$$\sum_{i \neq j} \varepsilon_{ij} J_{ij} \frac{\partial \zeta_{ij}}{\partial n_i} + 2q_j n_j = 0, \forall j, \quad (1a)$$

and $n_j \cdot n_j = 1, \forall j$ where the j is a constant in the summation.

$$\frac{\partial \zeta_{ij}}{\partial n_i} = \frac{n_j}{\sin(\zeta_{ij})}, \quad (2)$$

where the vector index has been suppressed on both the sides.

Using equation (2) in equation (1a) and taking the wedge product of the equation with n_j we have,

$$\sum_{i \neq j} \varepsilon_{ij} J_{ij} \frac{n_i \wedge n_j}{\sin(\zeta_{ij})} = 0, \forall j.$$

If

$$E_{ij} = \varepsilon_{ij} J_{ij} \frac{n_i \wedge n_j}{\sin(\zeta_{ij})},$$

then the last equation can be simplified to

$$\sum_{i \neq j} E_{ij} = 0, \forall j. \quad (3)$$

We now consider the properties of E_{ij} :

- Each i represents a tetrahedron. There are ten E_{ij} 's, each one of them is associated with one triangle of the four-simplex.
- The square of E_{ij} :

$$\begin{aligned} E_{ij} \cdot E_{ij} &= \frac{J_{ij}^2}{\sin^2(\zeta_{ij})} (n_j^2 n_i^2 - (n_i \cdot n_j)^2) \\ &= \frac{J_{ij}^2}{\sin^2(\zeta_{ij})} (1 - (\cos(\zeta_{ij}))^2) \\ &= J_{ij}^2. \end{aligned}$$

- The wedge product of any two E_{ij} is zero if they are equal to each other or if their corresponding triangles belong to the same tetrahedron.
- Sum of all the E_{ij} belonging to the same tetrahedron are zero according to equation (3).

It is clear that these properties contain the first four Barrett-Crane constraints[2]. So we have successfully extracted the bivectors corresponding to the triangles of a general flat four-simplex in Riemannian general relativity and the n_i are the normal vectors of the tetrahedra. The J_{ij} are the areas of the triangle as

one would expect. Since we did not impose any non-degeneracy Barrett-Crane conditions [2], it is not guaranteed that the tetrahedra or the four-simplex have non-zero volumes.

The asymptotic limit of the partition function of the entire simplicial manifold with triangulation Δ is

$$S(\Delta, \{n_{is} \in S^3, J_{ij}, \varepsilon_{ijs}\}) = \sum_{i < j, s} \varepsilon_{ijs} J_{ij} \zeta_{ijs},$$

where I have assumed variable s represents the four simplices of Δ and i, j represents the tetrahedra. The ε_{ijs} can be interpreted as the orientation of the triangles. Each triangle has a corresponding J_{ij} . The n_{is} denote the vector associated with the side of the tetrahedron i facing the inside of a simplex s . Now there is one bivector E_{sij} associated with each side facing inside of a simplex s of a triangle ij defined by

$$E_{ijs} = \varepsilon_{ijs} J_{ij} \frac{n_{js} \wedge n_{is}}{\sin(\zeta_{ijs})}.$$

If the n_{is} are chosen such that they satisfy stationary conditions

$$\sum_{i \neq j} E_{ijs} = 0, \forall j, s,$$

and if

$$\theta_{ij} = \sum_s \varepsilon_{ijs} \zeta_{ijs},$$

then

$$\begin{aligned} S(\Delta, \{J_{ij}, \varepsilon_{ijs}\}) &= \sum_{i < j, s} \varepsilon_{ijs} J_{ij} \zeta_{ijs}, \\ &= \sum_{i < j} J_{ij} \theta_{ij} \end{aligned}$$

can be considered to describe the Regge calculus [1] for the Riemannian general relativity. The angle θ_{ij} are the deficit angles associated with the triangles and the n_{is} are the vector normals associated with the tetrahedra.

References

- [1] Regge. T., General Relativity without Coordinates, Nuovo Cimento A 19, 558-571.
- [2] J. W. Barrett and L. Crane, Relativistic Spin Networks and Quantum general relativity. J.Math.Phys., 39:3296–3302, 1998.
- [3] J. W. Barrett and L. Crane, A Lorentzian Signature Model for Quantum General relativity , Class.Quant.Grav., 17:3101–3118, 2000.

- [4] J. W. Barrett and R. Williams, The asymptotics of an amplitude for the 4-Simplex, Preprint available as arXiv:gr-qc/9809032.
- [5] J. W. Barrett and C. Steele, Asymptotics of relativistic spin networks, Preprint available as arXiv:gr-qc/0209023.
- [6] G. Ponzano and T. Regge, Semiclassical Limit of Racah Coefficients, Spectroscopy and Group Theoretical Methods in Physics, F. Block et al (Eds), North-Holland, Amsterdam, 1968.
- [7] Klaus Schulten and Roy G. Gordon, Semiclassical approximations to 3j- and 6j- coefficients for quantum-mechanical coupling of angular momenta, J.Math.Phys, Vol.16, No.10, October 1975.